

For the last version of this text, type *hodgegaillard* on Google. Date of this version: Mon Dec 24 09:01:37 CET 2007.

A Hodge Theorem for Noncompact Manifolds

Theorem *If M is a riemannian manifold, then the inclusion of the complex of coclosed harmonic forms into the de Rham complex induces a linear isomorphism in cohomology. If M has at most countably many connected components, this linear isomorphism is a Fréchet isomorphism.*

[Manifolds are assumed to be C^∞ and Hausdorff.]

Theorem 5 in Section I.9.10 of Bourbaki [2] implying that M is paracompact, we can assume that it is connected, and also that it is non-compact (the result being classical in the compact case). Then the claim follows easily (using the Open Mapping Theorem and the fact that the de Rham cohomology is a Fréchet space) from the surjectivity of the laplacian on the de Rham complex. Let us check this surjectivity. In [4, p. 158] de Rham proves (using results of Aronszajn, Krzywicki and Szarski [1]) that a harmonic form which has a zero of infinite order vanishes identically; this implies in particular that the laplacian satisfies Property (A) in Definition 5 of Malgrange [3, p. 333]; it is well known that the laplacian satisfies also Condition (P) — called **ellipticity** nowadays — in Definition 6 of [3, p. 338]; in view of Theorem 5 in [3, p. 341] this implies the desired surjectivity.

- [1] Aronszajn N., Krzywicki A., Szarski J., A unique continuation theorem for exterior differential forms on Riemannian manifolds, *Ark. Mat.*, Volume 4, Number 5 (1962) 417-453. Available at springerlink.com.
- [2] Bourbaki, N., **Topologie générale**, Vol. 1, Chapitres 1 à 4, Hermann, 1971.
- [3] Malgrange B., Existence et approximation des solutions des équations aux dérivées partielles et des équations de convolution, *Ann. Inst. Fourier, Grenoble* **6** (1955-56) 271-354. Available at numdam.org.
- [4] de Rham G., **Differentiable manifolds**, Springer-Verlag, 1984.